

Dynamic depletion in a Bose condensate via a sudden increase of the scattering length

C. K. Law, P. T. Leung and M.-C. Chu

*Department of Physics, The Chinese University of Hong Kong,
Shatin, NT, Hong Kong, China*

(February 1, 2008)

We examine the time-dependent quantum depletion of a trapped Bose condensate arising from a rapid increase of the scattering length. Our solution indicates that a significant buildup of incoherent atoms can occur within a characteristic time short compared with the harmonic trap period. We discuss how the depletion density and the characteristic time depend on the physical parameters of the condensate.

PACS numbers: 03.75.Fi, 05.30.Jp

Bose-Einstein condensates (BEC) of atomic gases with tunable self-interaction strengths can be realized by applying strong magnetic fields near a Feshbach resonance [1,2]. Recent experiments have demonstrated dramatic dynamic features by manipulating the values of s-wave scattering lengths. For examples, a change of the interaction sign from repulsive to attractive can trigger a condensate collapse [3,4], a sweep of the magnetic fields across a resonance leads to an enhanced rate of inelastic collisions [5]. More recently Claussen *et al.* employed a magnetic field pulse to turn on a large scattering length in an extreme short time [6]. Their observation of condensate loss and its counterintuitive dependence on the pulse length suggest rich microscopic dynamics not captured by the mean field theory.

In this paper we address the interesting question how a sudden increase of the self-interaction strength affects the macroscopic quantum coherence. In particular we will examine the dynamic creation of incoherent atoms from the condensate (i.e., a time-dependent depletion effect) based on a two-body interaction model. Unlike the study of stationary systems [7], the calculation of depletion in nonstationary systems is a more difficult task because one needs to follow the time evolution of all collective excitation modes (assuming these modes are well defined in the remote past when the system is stationary) [8]. In this paper we present an approximate solution to tackle a time-varying scattering length problem. We find that the incoherent atom density can quickly build up if the change of scattering length is sufficiently large and rapid. Therefore a sudden change of interaction strength could lead to observable decoherence effects in the condensate. Our solution indicates the time scale of the depletion dynamics and the conditions that determine when the mean field theory breaks down.

To begin we consider a Bose-Einstein condensate of N

atoms confined in a spherical harmonic potential with a trap frequency ω_T . The time-dependence of the s-wave scattering length $a(t)$ is controlled by an external magnetic field pulse. We assume that the magnetic field does not cross any resonance, and possible molecule formation processes will not be considered. Under the diluteness assumption of the system $na(t)^3 \ll 1$ (n is the atom number density), the interaction between atoms is modelled by a two-body δ -potential. The second quantized time-dependent Hamiltonian of our model is given by,

$$H = \int d^3x \left(\hat{\Psi}^\dagger h_0 \hat{\Psi} + \frac{2\pi\hbar^2 a(t)}{m} \hat{\Psi}^\dagger \hat{\Psi}^\dagger \hat{\Psi} \hat{\Psi} \right) \quad (1)$$

Here $\hat{\Psi}$ is the atomic field operator, m is the atomic mass and $h_0 \equiv -\frac{\hbar^2}{2m} \nabla^2 + \frac{1}{2} m \omega_T^2 r^2$ corresponds to the trap Hamiltonian for a single particle. We note that the validity of replacing the real interatomic potential by the δ -potential in time dependent situations is a subtle issue. However, it is plausible that the δ -potential is valid if the rate of change of the system is small compare with a characteristic rate τ_s^{-1} that atoms need to adjust their relative wave functions [9]. The τ_s is typically a short time depending on the scattering length, we will discuss this issue later in this paper.

In this paper we assume that the condensate is stationary initially with a zero scattering length, i.e., $a(0) = 0$. Such an initial condition can be achieved by turning off the self-interaction of the condensate adiabatically via a magnetic field [6]. At later time, $a(t)$ is quickly raised to a peak value $\bar{a} > 0$. The rise time interval is sufficiently short (but longer than τ_s) such that the evolution of the condensate is negligible during the turn on. This allows us to approximate $a(t)$ in the form

$$a(t) \approx \bar{a} \theta(t), \quad (2)$$

where θ is the step function.

We decompose the field operator into a dominant coherent part and a small fluctuation part, i.e., $\hat{\Psi} = \sqrt{N} \Phi_0 + \hat{\psi}$. We require the condensate wave function Φ_0 to obey the time dependent mean field equation,

$$i\hbar \frac{\partial \Phi_0}{\partial t} = h_0 \Phi_0 + g(t) |\Phi_0|^2 \Phi_0, \quad (3)$$

where $g(t) \equiv 4\pi\hbar^2 a(t) N/m$. Then according to the Heisenberg equation of $\hat{\Psi}$, the fluctuation $\hat{\psi}$ obeys the equation,

$$i\hbar \frac{\partial \hat{\psi}}{\partial t} = \left(h_0 + 2g(t) |\Phi_0|^2 \right) \hat{\psi} + g(t) \Phi_0^2 \hat{\psi}^\dagger. \quad (4)$$

Note that in writing Eq. (4) we have assumed that $\hat{\psi}$ is small so that second and higher order terms of $\hat{\psi}$ are discarded.

Our task is to solve the time evolution of the operator $\hat{\psi}$ and examine the growth of incoherent atom density $\langle \hat{\psi}^\dagger(x, t) \hat{\psi}(x, t) \rangle$. First we need to specify the initial conditions for Φ_0 and $\hat{\psi}$. Since the system is noninteracting until $t = 0$, the initial condensate wave function Φ_0 is just the ground state of the harmonic trap. The corresponding quantum fluctuations at zero temperature is described by a vacuum state $|vac\rangle$ associated with $\hat{\psi}(\hat{x}, 0)$, i.e., $\hat{\psi}(\hat{x}, 0)|vac\rangle = 0$. Using a standard expansion in terms of the single particle eigenfunctions of the trap, the initial conditions are given by,

$$\Phi_0(\vec{x}, 0) = f_0(\vec{x}) \quad (5)$$

$$\hat{\psi}(\vec{x}, 0) = \sum_n \hat{b}_n f_n(\vec{x}). \quad (6)$$

Here $\hat{b}_n$ is the annihilation operator associated with the trap eigenfunction f_n obtained from $h_0 f_n = \hbar\omega_n f_n$, with f_0 being the lowest state. The initial state $|vac\rangle$ is the vacuum of $\hat{b}_n$.

The linear structure of Eq. (4) suggests a general solution in the form

$$\hat{\psi}(\vec{x}, t) = \sum_n \hat{b}_n u_n(\vec{x}, t) + \hat{b}_n^\dagger v_n^*(\vec{x}, t), \quad (7)$$

where the $\hat{b}_n$ and $\hat{b}_n^\dagger$ are time-independent operators defined in Eq. (6). A direct substitution of (7) in (4) gives the equation of motion of the mode functions $u_n(\vec{x}, t)$ and $v_n(\vec{x}, t)$:

$$i\hbar \frac{\partial u_n}{\partial t} = \left(h_0 + 2g(t) |\Phi_0(\vec{x}, t)|^2 \right) u_n + g(t) \Phi_0^2(\vec{x}, t) v_n \quad (8)$$

$$i\hbar \frac{\partial v_n}{\partial t} = - \left(h_0 + 2g(t) |\Phi_0(\vec{x}, t)|^2 \right) v_n - g(t) \Phi_0^{*2}(\vec{x}, t) u_n. \quad (9)$$

To match the initial condition (6), we set

$$u_n(\vec{x}, 0) = f_n(\vec{x}) \quad (10)$$

$$v_n(\vec{x}, 0) = 0. \quad (11)$$

Thus the operator equation (4) with the initial condition (6) can now be replaced by the c -number equations (8-11). For later purpose, it is convenient to introduce the quantity $\lambda(\vec{x}) \equiv 4\pi\hbar a N f_0^2(\vec{x})/m$ that characterizes the position dependent self-interaction in frequency unit.

In this paper we present a solution in the short time domain $0 \leq t \leq T_0$ in which T_0 is defined by $T_0 \equiv 1/\lambda(0)$ with $\omega_T T_0 \ll 1$. In this domain the condensate wave function does not have enough time to evolve spatially. The condensate can only pick up a self-modulated phase arising from the self-interaction, i.e.,

$$\Phi_0(\vec{x}, t) \approx f_0(\vec{x}) e^{-i\lambda(\vec{x})t}. \quad (12)$$

Note that the trivial phase factor $e^{-i\omega_T t/2}$ due to the trap ground state energy is ignored since $\omega_T t \ll 1$ under the stated conditions above. We can easily verify the solution by a substitution in Eq. (3). The approximation we made is the discard of terms involving the spatial gradient of $e^{-i\lambda(\vec{x})t}$. These terms would describe the expansion of the condensate but they only become significant in a longer time scale $\approx 1/\sqrt{\omega_T \lambda(0)} > T_0$, which is beyond our specified time domain. We have also performed numerical tests that show Eq. (12) has good agreement with the exact numerical solution of Eq. (3) within the specified time domain.

We can solve Eq. (8) and (9) in a similar fashion. We write

$$u_n(\vec{x}, t) = f_n(\vec{x}) U_n(\vec{x}, t) + \delta_{nu}(\vec{x}, t) \quad (13)$$

$$v_n(\vec{x}, t) = f_n(\vec{x}) V_n(\vec{x}, t) + \delta_{nv}(\vec{x}, t) \quad (14)$$

where the functions U_n and V_n are defined by,

$$i\hbar \frac{\partial U_n}{\partial t} = \left(\hbar\omega_n + 2g(t) |\Phi_0(\vec{x}, t)|^2 \right) U_n + g(t) \Phi_0^2(\vec{x}, t) V_n \quad (15)$$

$$i\hbar \frac{\partial V_n}{\partial t} = - \left(\hbar\omega_n + 2g(t) |\Phi_0(\vec{x}, t)|^2 \right) V_n - g(t) \Phi_0^{*2}(\vec{x}, t) U_n \quad (16)$$

subjected to the condition $U_n(\vec{x}, 0) = 1$ and $V_n(\vec{x}, 0) = 0$. We shall show that the δ_{nu} and δ_{nv} are small correction terms in the short time domain that we are concerned.

Let us write down the explicit form of U_n and V_n . Since Eq. (15) and (16) do not involve any gradient operator, different spatial coordinates are decoupled. At each position the equations behave as a two-level system that can be exactly solved:

$$U_n(\vec{x}, t) = e^{-i\lambda(\vec{x})t} \left[\cos \Omega_n(\vec{x})t - i \frac{\omega_n + \lambda(\vec{x})}{\Omega_n(\vec{x})} \sin \Omega_n(\vec{x})t \right] \quad (17)$$

$$V_n(\vec{x}, t) = i e^{i\lambda(\vec{x})t} \frac{\lambda(\vec{x})}{\Omega_n(\vec{x})} \sin \Omega_n(\vec{x})t \quad (18)$$

which satisfy the normalization $|U_n|^2 - |V_n|^2 = 1$, and the quantity

$$\Omega_n(\vec{x}) = \sqrt{\omega_n (\omega_n + 2\lambda(\vec{x}))} \quad (19)$$

can be interpreted as a form of frequency analogous to the Rabi oscillations in two-level systems.

To show δ_{nu} and δ_{nv} are small, we note that they are both zero initially. Their equations of motion can be obtained by a direct substitution of Eq. (13) and (14) into (8) and (9). This leads to a set of linear equations similar to Eq. (8) and (9) but with inhomogeneous terms as

$$i \frac{\partial \delta_{nu}(\vec{x}, t)}{\partial t} = -\frac{\hbar}{m} \left(\nabla U_n \cdot \nabla f_n - \frac{1}{2} f_n \nabla^2 U_n \right) + \dots \quad (20)$$

$$i \frac{\partial \delta_{nv}(\vec{x}, t)}{\partial t} = \frac{\hbar}{m} \left(\nabla V_n \cdot \nabla f_n - \frac{1}{2} f_n \nabla^2 V_n \right) + \dots \quad (21)$$

These inhomogeneous terms involve the spatial gradient of f_n , U_n and V_n . Since U_n and V_n are smooth functions in the length scale of the trap ground state, the dominant contribution comes from ∇f_n for high excited modes. We can estimate the size of δ_{nv} using the first order approximation which is a time integration of the inhomogeneous term $\nabla V_n \cdot \nabla f_n$. Up to the time $t = T_0$, we find that δ_{nv} is only a small fraction of $f_n V_n$. The fraction is typically of the order $(\omega_T T_0)^{1/2} \ll 1$ [10]. A similar conclusion applies to δ_{nu} .

The functions $f_n U_n$ and $f_n V_n$ are approximate solutions of u_n and v_n . The approximation comes from the fact that $\nabla^2 (f_n U_n) \approx U_n \nabla^2 f_n$ and $\nabla^2 (f_n V_n) \approx V_n \nabla^2 f_n$ are the leading contributions in the Laplacians for slowly varying (spatially) functions U_n and V_n . The correction terms can remain small under the conditions stated in the previous paragraph. We have tested the approximate scheme beyond the first order consideration by solving Eq. (8) and (9) numerically in a one-dimensional model, the exact results are in good agreement with those obtained by the approximate method.

Now we come to the main result of our paper. The time-development of incoherent atom density after the change of the scattering length is given by,

$$\begin{aligned} \langle \hat{\psi}^\dagger(\vec{x}, t) \hat{\psi}(\vec{x}, t) \rangle &= \sum_n |v_n(\vec{x}, t)|^2 \\ &\approx \sum_n |f_n(\vec{x})|^2 |V_n(\vec{x}, t)|^2. \end{aligned} \quad (22)$$

With the help of Eq. (18), we have,

$$\begin{aligned} \langle \hat{\psi}^\dagger(\vec{x}, t) \hat{\psi}(\vec{x}, t) \rangle &\approx \sum_n |f_n(\vec{x})|^2 \frac{\lambda^2(\vec{x})}{\omega_n(\omega_n + 2\lambda(\vec{x}))} \\ &\times \sin^2 \left[\sqrt{\omega_n(\omega_n + 2\lambda(\vec{x}))} t \right]. \end{aligned} \quad (23)$$

For a spherical harmonic trap, each single particle state n is described by three quantum numbers (n_1, n_2, n_3) in Cartesian coordinates, and the corresponding frequency $\omega_n = (n_1 + n_2 + n_3 + 3/2)\omega_T$. Therefore Eq. (23) is a triple summation.

Let us calculate the incoherent atom density at the trap center $\vec{x} = 0$. It is a good approximation (particularly for high modes) that

$$\begin{aligned} |f_{2j_1, 2j_2, 2j_3}(0)|^2 &\approx 0.09 \times l_0^{-3} \\ &\times \left[\left(2j_1 + \frac{1}{2} \right) \left(2j_2 + \frac{1}{2} \right) \left(2j_3 + \frac{1}{2} \right) \right]^{-1/2} \end{aligned} \quad (24)$$

where $l_0 = \sqrt{\hbar/m\omega_T}$ and we need only to consider even modes at the origin. Eq. (24) is obtained from the semiclassical construction of the trap eigenfunctions. One

can check that Eq. (24) is fairly accurate even for the ground state. Next we change the variables j_i by introducing: $\tilde{k}_i^2 = (4j_i + 1)\omega_T T_0/2$ for $i = 1, 2, 3$. We find that the function inside the summation (23) changes smoothly with $\tilde{k}_i$. Therefore we may replace the discrete sum (23) by an integral that can further be simplified using the spherical symmetry. In order to compare the fluctuations with the mean-field density, we define $\Delta(t) \equiv \langle \hat{\psi}^\dagger(0, t) \hat{\psi}(0, t) \rangle / n_0$ which is the ratio of the central incoherent atom density to the condensate density $n_0 = N f_0^2(0)$. We find that

$$\Delta(t) \approx 6.3(n_0 \bar{a}^3)^{1/2} \int_0^\infty dk \frac{\sin^2(kt\sqrt{2+k^2}/T_0)}{2+k^2} \quad (25)$$

where $k^2 = \tilde{k}_1^2 + \tilde{k}_2^2 + \tilde{k}_3^2$. The integral in Eq. (25) is an increasing function of time, but it must be less than $\pi/2\sqrt{2} \approx 1.1$. At the time $t = T_0$ the integral reaches 0.53 that gives $\Delta \approx 3.4(n_0 \bar{a}^3)^{1/2}$ [11]. This result is about twice the value that one would obtain from a (stationary) homogeneous condensate at zero temperature [12]. Note that $\Delta(t)$ depends explicitly on the gas parameter $n_0 \bar{a}^3$ in Eq. (25). If $\Delta(t)$ is comparable to one, then both the mean field theory and the linearized quantum theory break down because quantum fluctuations are no longer small. For a general position $\vec{x}$, we can work out a similar integral. The spatial profile can be well fitted by a Gaussian: $\langle \hat{\psi}^\dagger(\vec{x}, t) \hat{\psi}(\vec{x}, t) \rangle \approx \langle \hat{\psi}^\dagger(0, t) \hat{\psi}(0, t) \rangle e^{-3r^2/2l_0^2}$.

Finally we would like to discuss the validity of the model (1). A necessary condition that the usual pseudopotential approach can be generalized to time-dependent situations is that the time-varying parameters of the system do not excite high relative wave vectors between atoms. This condition ensures $pa/\hbar \ll 1$ (where p is the relative momentum) required for the s-wave approximation. There is no simple way to check the condition for general interatomic potentials, but it is useful to consider the hard sphere model as an illustrative example. In the relative coordinate of two hard spheres, the expanding spheres are described by a moving boundary condition. If the radial velocity is $\dot{a}$, a slow incoming wave will pick up a wavevector of magnitude roughly $m_r \dot{a}/\hbar$ by reflection, where $m_r = m/2$ is the reduced mass. Therefore $m_r \dot{a}a/\hbar \ll 1$ is the required condition that specifies how slow the speed should be. If we consider $a(t) = \dot{a}t/t_{rise}$ is linear in time during a rise period t_{rise} , then the condition is satisfied if $m_r \dot{a}^2/\hbar \ll t_{rise}$. Quite remarkably, the characteristic time $\tau_s \equiv m_r \bar{a}^2/\hbar$ estimated here agrees with the one suggested by Leggett under a general consideration [9].

To give a physical example, we consider a ^{85}Rb condensate with 10^5 atoms in a spherical trap with $\omega_T = 2\pi \times 10$ Hz. If the scattering length $\bar{a} = 30$ nm, then $T_0 = 80\mu\text{s}$ and $\tau_s = 1\mu\text{s}$ is a short time. In this example, the gas parameter $n_0 \bar{a}^3 = 0.001$ and $\Delta = 0.11$ at $t = T_0$. This means a 11% depletion of condensate density at the trap

center in $80\mu s$. We may compare this example with the case with a smaller scattering length $\bar{a} = 3$ nm (same particle number and trap frequency) that only gives 0.36% depletion density in $800\mu s$.

In conclusion we made use of the model (1) to analyze the dynamic depletion in a Bose condensate after a sudden increase of the self-interaction strength. The depletion is a quantum decoherence process originated from the vacuum fluctuations associated with $\hat{\psi}$. Our solution (23) and (25) indicate how the depletion density increases as a function of time. We discover that $T_0 = m/4\pi\hbar\bar{a}Nf_0^2(0)$ is an important time scale for the buildup of incoherent atoms for a trapped condensate with $\omega_T T_0 \ll 1$. It is interesting to rewrite T_0 in the form: $T_0 = \omega_T^{-1}(n_0\bar{a}^3)^{-1/3}(n_0l_0^3)^{-2/3}/4\pi$. This indicates the relationship between the characteristic time T_0 and different parameters, including the gas parameter. Finally, we point out that the recent experiment at JILA [6] reaching $n_0\bar{a}^3 \approx 0.3$ is beyond the scope of the present theory. Such a strong coupling would give $\tau_s \approx 39\mu s$ that is *longer* than the rise time of the magnetic pulse, and so the s-wave approximation model is questionable. The microscopic processes in such an extreme situation remain an interesting open topic for investigations.

ACKNOWLEDGMENTS

We acknowledge the support from an RGC Earnmarked Grant CUHK4237/01P and the Chinese University of Hong Kong Direct Grant (Grant Nos: 2060148 and 2060150).

$\Delta(2T_0) \approx 3.5(n_0a^3)^{1/2}$ which is only about a 3% increase compared with the value at $t = T_0$.

[12] K. Huang, *Statistical Mechanics* (Wiley, New York 1987).

-
- [1] S. Inouye *et al.*, Nature **392**, 151 (1998).
 - [2] S. L. Cornish *et al.*, Phys. Rev. Lett. **85**, 1795 (2000).
 - [3] J. L. Roberts *et al.*, Phys. Rev. Lett. **86**, 4211 (2001).
 - [4] E. A. Donley *et al.*, Nature **412**, 295 (2001).
 - [5] J. Stenger *et al.*, Phys. Rev. Lett. **82**, 2422 (1999).
 - [6] N. R. Claussen *et al.*, cond-mat/0201400.
 - [7] J. Javanainen, Phys. Rev. A **54**, 3722 (1996); K. G. Singh and D. S. Rokhsar, Phys. Rev. Lett. **77**, 1667 (1996); F. Dalfovo *et al.*, Phys. Rev. A **56**, 3840 (1997).
 - [8] Y. Castin and R. Dum, Phys. Rev. Lett. **79**, 3553 (1997).
 - [9] A. J. Leggett, Rev. Mod. Phys. **73**, 307 (2001).
 - [10] Our approximation works even better at $r = 0$, since $\nabla V_n \cdot \nabla f_n$ and $\nabla U_n \cdot \nabla f_n$ both vanish at the center. The non-zero inhomogeneous terms $f_n \nabla^2 V_n$ and $f_n \nabla^2 U_n$ contribute a correction of order $\omega_T T_0$ instead of $(\omega_T T_0)^{1/2}$.
 - [11] The growth of Δ is most significant within the time T_0 after the interaction is turned on. Extending the duration does not yield a much increase of Δ . For example, for the same condensate, twice the duration (i.e., $2T_0$) gives